



CONDITION	EXCLUSIONS TO TREATMENT
<b>ACNE</b> (MILD TO MODERATE)	Under 13 years old or over 65 years old Severe acne Eczematous, broken or sunburnt skin
<b>ATHLETES FOOT</b>	Pregnant or breastfeeding Diabetes Toenails are black or discoloured
<b>CONJUNCTIVITIS</b> (SUPERFICIAL EYE INFECTIONS)	Child under 3 months old Severe infection Recent course of chloramphenicol Eye pain or injury to eye Eye surgery in last six months Glaucoma
<b>CONSTIPATION</b>	Children under 5 years old if a recurring problem Abdominal pain, vomiting, bloating or pain on defecation
<b>CYSTITIS</b> (UNCOMPLICATED URINARY TRACT INFECTION IN WOMEN)	Under 18 years or over 65 years old Pregnant or breastfeeding Fever or feeling unwell Confused or dehydrated Treatment within previous three months
<b>DIARRHOEA</b>	Children under 6 months Significant weight loss Diabetes Presence of blood or mucus in stools Bile stained vomiting
<b>GOUT</b>	Under 18 years old No previous diagnosis Frequent attacks
<b>HAYFEVER</b> (SEASONAL ALLERGIC RHINITIS)	Pregnant or breastfeeding Thick yellow or green discharge from eyes or nose Persistent or painful ear or sinusitis
<b>HEAD LICE</b> (MUST BE LIVE LICE)	None
<b>INDIGESTION / HEARTBURN</b>	Children under 12 years old Pregnant or breastfeeding Blood in stools Persistent vomiting Significant weight loss Previous history of or suspected duodenal ulcer/ gastric ulcer
<b>IMPETIGO</b> (SMALL AREA)	Children under 1 year Pregnant or breastfeeding Currently taking a prescribed antibiotic More than 2 small areas and each area no greater than the size of a one pence piece Any course of treatment for Impetigo within 6 months

CONDITION	EXCLUSIONS TO TREATMENT
<b>MIGRAINE</b>	No previous diagnosis Under 18 years or over 65 years old Pregnant or breastfeeding Prophylactic treatment Experiencing any of the following symptoms: double vision, clumsy and uncoordinated movements, tinnitus or recent onset of rash with headache
<b>ORAL THRUSH</b> (ORAL CANDIDIASIS) <b>IN INFANTS</b>  AND IF NEEDED; TREATMENT OF <b>CANDIDA (THRUSH) OF THE NIPPLE</b> <b>IN BREASTFEEDING MOTHERS</b>	Babies under 1 month old or over 12 months old Child is hypersensitive or allergic to nystatin Child is unwell Reoccurrence within last 8 weeks  Fever, temperature and feeling unwell Pregnant This episode is a recurrence within last 8 weeks
<b>PILES</b> (HAEMORRHOIDS)	Under 18 years old Pregnant or breastfeeding Bleeding Chronic irritation or leaking Abdominal pain/vomiting Patients taking anticoagulants (eg warfarin)
<b>RASH</b> (MILD DERMATITIS/ECZEMA)	Face or genital area Contact dermatitis or infected eczema Severe, widespread and/or oozing Caused by medication Nappy rash
<b>SCABIES</b>	Children under 2 years old Significant irritation and/or inflammation Crusted scabies
<b>THREADWORM</b>	Children under 2 years old Pregnant or breastfeeding Tummy pain or rectal bleeding (bleeding from the bottom) Significant, unexplained weight loss Significant change in bowel habit Taking metronidazole or cimetidine
<b>VAGINAL THRUSH</b>	Women under 16 years or over 60 years old Pregnant Previous treatment within 3-4 weeks or twice in 6 months Diabetes Abnormal menstrual bleed Suspected sexually transmitted infection First occurrence of symptoms

